

Material-barrier Tunneling in One-dimensional Few-boson Mixtures

Anika C. Pflanze¹, Sascha Zöllner^{2,*} and Peter Schmelcher^{1,3,†}

¹*Physikalisches Institut, Universität Heidelberg,
Philosophenweg 12, 69120 Heidelberg, Germany*

²*Niels Bohr International Academy, Niels Bohr Institute,
Blegdamsvej 17, 2100 København, Denmark*

³*Theoretische Chemie, Universität Heidelberg,
Im Neuenheimer Feld 229, 69120 Heidelberg, Germany*

(Dated: September 9, 2009)

Abstract

We study the quantum dynamics of strongly interacting few-boson mixtures in one-dimensional traps. If one species is strongly localized compared to the other (e.g., much heavier), it can serve as an effective potential barrier for that mobile component. Near the limit of infinite localization, we map this to a system of identical bosons in a double well. For realistic localization, the backaction of the light species on the “barrier” atoms is explained—to lowest order—in terms of an induced attraction between these. Even in equilibrium, this may outweigh the bare intra-species interaction, leading to unexpected correlated states. Remarkably, the backaction drastically affects the inter-species dynamics, such as the tunneling of an attractively bound pair of fermionized atoms.

PACS numbers: 03.75.Mn, 03.65.Xp, 03.75.Lm

*Electronic address: zoellner@nbi.dk

†Electronic address: peter.schmelcher@pci.uni-heidelberg.de

Tunneling of particles through an energetically forbidden region is a hallmark quantum-mechanical effect, which illustrates the wave nature of matter. The experimental flexibility of ultracold atoms makes it possible to study this phenomenon in a clean and highly controlled environment, where the role of the tunnel barrier is played, e.g., by light forces such as in optical lattices [1]. This has fostered the direct observation of fundamental effects like second-order tunneling [2], Josephson oscillations and nonlinear self-trapping [3]. Moreover, optical lattices have proven powerful quantum simulators, giving insight into the role of tunneling in, e.g., the quantum phase transition from superfluid to insulator [4], or in spin-exchange processes responsible for quantum magnetism [5].

All of these cases reflect the paradigm of tunneling through a *classical* barrier described by some external potential. In this Letter, we investigate tunneling through a material or *quantum* barrier, in the sense that it interacts with the particles. This is realized via quasi-one-dimensional (1D) mixtures of two atomic species, one of which is squeezed in the trap center. When the localization is very tight, we show that this can indeed be understood as an effective tunnel barrier for the other species. As that confinement is relaxed, the “barrier” atoms move due to the backaction of the other species. The dramatic effect of this correlation on the mobile species is studied both for the ground state and for the inter-species tunneling dynamics. To first explore its microscopic mechanism, we consider a few-atom system, which is studied in a numerically exact fashion utilizing the multi-configuration time-dependent Hartree method [6]. This wave-packet dynamics tool which has been applied successfully to systems of identical bosons as well as mixtures (see [7] for details). We then go on to derive approximate models that capture the relevant physics for higher atom numbers. Still, such a small system may be achieved experimentally, e.g., by creating arrays of 1D optical lattices (such as in [5]) or of 1D tubes [8], each containing only a few atoms. Moreover, high-resolution imaging techniques such as scanning-electron microscopy allow for single-site addressability [9].

Model.— We consider a mixture of two 1D bosonic species, labeled $\sigma = A, B$. These may correspond to different atomic species (or isotopes); however, in the case of equal masses they can also be thought of as different hyperfine components. The many-body Hamiltonian then reads $H = \sum_{\sigma} H_{\sigma} + H_{AB}$, with the single-species Hamiltonian $H_{\sigma} = \sum_{i=1}^{N_{\sigma}} \left[\frac{1}{2m_{\sigma}} p_{\sigma,i}^2 + U_{\sigma}(x_{\sigma,i}) \right] + \sum_{i < j} g_{\sigma} \delta(x_{\sigma,i} - x_{\sigma,j})$ and the inter-species coupling $H_{AB} = \sum_{i,j} g_{AB} \delta(x_{A,i} - x_{B,j})$. In what follows, we will focus on the case of harmonic trapping potentials $U_{\sigma}(x) = \frac{1}{2} m_{\sigma} \omega_{\sigma}^2 x^2$ and repulsive forces, $g_{\sigma}, g_{AB} \geq 0$. (Note that taking $g_{\sigma} \rightarrow \infty$, the component σ can be mapped to a fermionic one

[10], referred to as *fermionization*. In this sense, considering bosonic atoms poses no restriction in 1D.) Rescaling to harmonic-oscillator units, we can eliminate $\hbar = m_A = \omega_A = 1$ [7].

Material tunneling barrier.— Imagine the situation where one of the species (say B) is much more strongly localized in the trap center. This can be achieved by drastically reducing its (unperturbed) length scale $a_B \equiv 1/\sqrt{m_B\omega_B} \ll 1$, which amounts to having a near-zero mass ratio $\alpha \equiv m_A/m_B \ll 1$ (for different atom masses such as in Li/Cs) and/or strong confinement of B, $\omega_A/\omega_B \ll 1$ (e.g., realized via species-dependent optical lattices [12]). For simplicity, let us assume much heavier B atoms but equal frequencies. This is not crucial, though, and experimentally a large frequency ω_B may enhance the localization effect.

Being tightly localized, the B atoms should feel no density variations of A and, to lowest order, the total density matrix $\hat{\rho}_{AB}^{(N)} \approx \hat{\rho}_A^{(N_A)} \otimes \hat{\rho}_B^{(N_B)}$ can be approximately factorized. (This even holds for strong repulsion $g_{AB} \gg 1$, since the strong localization of the B atoms at the trap center effectively reduces the interaction to a single-particle potential [14].) Integrating out the heavy B atoms leads to an effective Hamiltonian for the light species [7],

$$\bar{H}_A^{(0)} = H_A + \text{tr}_B[H_{AB}\hat{\rho}_B^{(N_B)}] = H_A + g_{AB}\sum_i n_B(x_{A,i}). \quad (1)$$

In this light, A ought to feel only an effective single-particle potential $g_{AB}n_B(x)$, which in our case likens a sharp barrier at $x = 0$, with a width $\sim a_B$ given by the one-body density $\rho_B \equiv n_B/N_B$ (in terms of the number density n_B) and its height proportional to the inter-species coupling g_{AB} . In particular, for $N_B g_{AB} \gg 1$, this barrier practically splits the harmonic trap for the A atoms into a pronounced double well. (By extension, localizing the B atoms over several centers should make for an effective “lattice” for A.) This scenario naturally brings up the question whether one can create an effective localization of the light atoms (in equilibrium) or even see dynamic effects such as tunneling solely due to inter-species interactions.

Figure 1 shows that, near the limit of strong localization, the simple picture invoked above is indeed consistent with our exact results. In detail, we have evidenced this on the ground state of $N_A = 4$ atoms in the presence of one localized “impurity” ($\alpha = 0.001$). Absent any intra-species interactions ($g_A = 0$), the A atoms are coherently spread over the left- and right-hand side of the trap in analogy to an external double-well potential [11], even though the system as a whole is strongly interacting ($g_{AB} = 25$ unless otherwise stated). This materializes in the two-body density $\rho_A^{(2)}(x_1, x_2)$, measuring the joint probability density of finding two A atoms at x_1 and x_2 : In Fig. 1(a), both are equally likely to be on the same site (peaks on the diagonal, $x_1 \approx x_2$) as on




Figure 1: (color online) Strong-localization limit, $\alpha \equiv m_A/m_B = 0.001$, for $N_A = 4$ atoms in the presence of a B atom localized at the center. *Top*: Two-body density $\rho_A^{(2)}(x_1, x_2)$ for $g_A = 0, 0.5$, and 25 from left to right. *Bottom*: Density profile $\rho_A(x)$ for different g_A . (All quantities in harmonic-oscillator units of A, see text.)

opposite ones ($x_1 \approx -x_2$). As g_A increases, the A atoms assume a “Mott-insulator”-type state with all A atoms more or less localized in either well [$g_A = 0.5$; Fig. 1(b)], and eventually *fermionize* [$g_A = 25$; Fig. 1(c)]. The checkerboard pattern emerging in the latter case can be understood simply as a gas of A-component bosons with hard-core repulsion (or, equivalently, noninteracting fermions) immersed in a *double-well* trap [11, 13]: In each well, measuring one A-boson at x_1 pins down the position of the remaining bosons to $N_A - 1$ discrete spots. This reflects in the density profile $\rho_A(x)$, where N_A marked density maxima form as $g_A \rightarrow \infty$ (Fig. 1, bottom).




Figure 2: (color online) Beyond the static-barrier picture: Two-body density $\rho_A^{(2)}(x_1, x_2)$ for (top) $g_A = 0$, with increasing mass ratios $\alpha = 0.001, 0.002, 0.02$ (from left to right); (bottom) $g_A = 0.5$, $\alpha = 0.001, 0.12, 0.2$. Same atom numbers as in Fig. 1.

Beyond the static limit.— We have so far proceeded on the assumption that the B atoms were heavy enough to be frozen out completely, that is, treated as a classical potential. For less restrictive mass ratios, however, these are expected to move due to the backaction of the light atoms. The impact of this correlation becomes palpable in Fig. 2, where the ground-state evolution of $\rho_A^{(2)}(x_1, x_2)$ is displayed as the mass ratio α is increased. For $g_A = 0$ (Fig. 2, top), the previously uncorrelated pattern develops into a strongly localized one ($\alpha = 0.02$), where all A atoms are found exclusively on the *same* site; i.e., they cluster. By contrast, the anti-correlated “insulating” ground state observed at $g_A = 0.5$ turns into a seemingly *uncorrelated* one as the B atom becomes less heavy (bottom).

To understand this wealth of phenomena, it would be desirable to extend the effective Hamiltonian (1) to higher orders in $\alpha > 0$. This is nontrivial since $H(\alpha)$ in itself does not suggest a straightforward power series in α . Our basic procedure is sketched in what follows (details will be given elsewhere [14]). Given the strong localization of the B atom ($N_B = 1$ without loss of generality), we introduce center-of-mass and Jacobian coordinates relative to $x_B \approx 0$. The resulting Hamiltonian affords an expansion in powers of α , which we can exploit so as

to derive an effective Hamiltonian for the A-atoms by tracing out B, which leaves us with $\bar{H}_A(\alpha) = \bar{H}_A^{(0)} + \alpha g_{AB} \frac{1}{2} \sum_{k \neq l} [(x_{A,k} - x_{A,l}) n'_B(x_{A,l}) - n_B(x_{A,l})] + O(\alpha^2)$. The first term recovers the initial infinite-mass approximation; the first-order correction in turn may be understood as an additional external potential $\delta U_A(x) \equiv -\alpha g_{AB} \frac{N_A-1}{2} [x n'_B(x) + n_B(x)]$, plus an induced (nonlocal) interaction between two A atoms: $\delta V_A(x_1, x_2) \equiv \alpha g_{AB} [x_1 n'_B(x_2) + x_2 n'_B(x_1)] / 2$, which adds to the local one $V_A(x_1, x_2) = g_A \delta(x_1 - x_2)$.

Given the effective multi-well (here: two sites) geometry experienced by the A atoms, it is tempting to explain some qualitative features from the perspective of simplified lattice models. Since, for $\alpha \rightarrow 0$, the single-particle physics of species A is governed by $\bar{h}_A \equiv \frac{1}{2} p^2 + \frac{1}{2} x^2 + g_{AB} n_B(x)$, it is natural to use its Bloch-type eigenstates (which for a double well are delocalized anti-/symmetric functions that come in bands of doublets) as a basis for the many-body Hamiltonian. In the spirit of the (bosonic) Hubbard model [15], it is actually preferable to introduce Wannier-type functions $w_s^{(\beta)}$ localized on the left (right) site s in band $\beta = 0, 1, \dots$. Expanding the many-body Hamiltonian $\bar{H}_A(\alpha)$ in terms of these Wannier functions and—in the case of sufficiently small interaction energies and deep wells—retaining only lowest-band and on-site terms yields an effective Bose-Hubbard model

$$\bar{H}_A^{(\text{BH})} = -J \sum_{\langle s, s' \rangle} \hat{a}_s^\dagger \hat{a}_{s'} + \frac{u}{2} \sum_s \hat{n}_s (\hat{n}_s - 1). \quad (2)$$

Here $-J = \langle w_s^{(0)} | \bar{h}_A + \delta U_A(x) | w_{s'}^{(0)} \rangle$ is the renormalized tunnel coupling of the lowest band, and $u = \langle w_s^{(0)\otimes 2} | V_A + \delta V_A | w_s^{(0)\otimes 2} \rangle = u_0 + \delta u$ denotes the on-site interaction. While the correction to J tends to be small, the on-site renormalization due to δV_A can have a huge effect if it is on the same order as the $\alpha = 0$ term. A closer analysis [14] reveals that $\delta u < 0$ always, signifying an *attractive* induced interaction. There is an intuitive way of picturing this self-interaction: If an A atom sits in one well, then it will repel the barrier; thus the well becomes more spacious, making it energetically favorable to accommodate yet another A atom.

Taking the induced attraction at face value, this casts a light on our results above. For $g_A = 0$ (Fig. 2, top), $u = \delta u < 0$ for a finite mass of B. This makes it plausible that the A atoms tend to cluster on *same* sites at $\alpha \sim 0.02$, although being really noninteracting. Likewise, for small repulsion as in Fig. 2 (bottom; $g_A = 0.5$), both terms may even cancel, $u_0 + \delta u \approx 0$ – this helps explain the seemingly uncorrelated pattern as α is increased.

Although giving the right trend for weak interactions between the A atoms, the validity of (2) is actually much more limited than that of the effective Hamiltonian $\bar{H}_A(\alpha)$. In particular, the Bose-




Figure 3: (color online) Inter-species tunneling dynamics of $N_A = 2$ bosons through a B atom: Relative A population of the right-hand side over time, $p_R(t)$, for different mass ratios, $\alpha = 0.001$ (—), $\alpha = 0.01$ (---), and $\alpha = 0.12$ (···). (a) $g_A = 0$, (b) $g_A = 0.5$, (c) $g_A = 25$; all $g_{AB} = 8$. *Insets*: Probability $p_2(t)$ of finding two A atoms on the same side.

Hubbard model breaks down for strong interactions comparable to the band gap. In that regime, rich multi-band effects can be found for stronger intra-species correlations, which we will discuss below in the context of the atoms' quantum dynamics.

Inter-species tunnel dynamics.— We have so far investigated the equilibrium situation of species A in the presence of an effective “barrier” composed of a second, localized component. It would be thrilling to learn how this affects the quantum dynamics – for instance, can the light atoms tunnel through the heavy ones, and how does the barrier atoms' motion influence this inter-species tunnel dynamics?

To answer this question, let us investigate the time evolution of the A bosons loaded initially in, say, the left-hand side of the trap, with the barrier atoms (B) tightly centered. This can be done, e.g., by displacing the trap center of $U_A(x)$ or by blocking the other half with a laser beam. Upon release, the A atoms may tunnel through species B, this way moving the barrier atoms, which in turn modifies the effective potential. To monitor the dynamics, we have recorded the percentage of A-atoms on the right, $p_R(t) = \int_0^\infty \rho_A(x;t)dx$, in dependence of α .

Figure 3(a) displays this population dynamics for $N_A = 2$ *noninteracting* A bosons ($g_A = 0$). For a static double well, we simply expect a Rabi-type oscillation of the population between left and right [16]. Indeed, for $\alpha = 0.001$, this is what we find. Increasing the mass ratio, the clear sine mode gives way to a more complex, two-mode oscillation ($\alpha = 0.01$), until for much larger values ($\alpha = 0.12$) the tunneling slows down drastically, with only a tiny faster modulation on top. This is reminiscent of second-order tunneling well known from repulsive atom pairs [2, 16]. In fact, the effective Bose-Hubbard model (2) suggests that this corresponds to *attractively* bound pairs in a double well, where single-atom Rabi tunneling is highly suppressed [17]. This line of reasoning is supported by the *pair (or same-site) probability* $p_2(t) = \int_{\{x_1, x_2 \geq 0\}} \rho_A^{(2)}(x_1, x_2; t) dx_1 dx_2$, measuring how likely it is to find two A-atoms on the same site [16]: This is plotted in the inset of Fig. 3(a), indicating that the atom pair becomes more and more stable (or p_2 no longer drops far below 1) as α is increased.

This contrasts with the case of $g_A = 0.5$, shown in Fig. 3(b). Near the static limit ($\alpha \leq 0.01$), one recovers the situation of repulsively bound pairs [2, 16], which tunnel at a period $T/2\pi \sim u/4J^2$ long compared with the Rabi oscillations $1/2J$, and which are stable in time (inset). Allowing for a finite mass of B, this pair breaks up, and the dynamics starts to resemble sinusoidal Rabi oscillations for $\alpha = 0.12$. This becomes even more conclusive from the angle of the pair probability (inset), whose minimum value now significantly deviates from unity. Qualitatively, this feature is captured by the effective Bose-Hubbard model (2): At large enough α , the attractive on-site interaction $\delta u < 0$ tends to cancel u_0 , which is in agreement with the Rabi-like oscillations observed here.

What happens for increasing intra-species repulsion, in particular upon approaching the fermionization limit, e.g., at $g_A = 25$ [Fig. 3(c)]? In the case of a quasi-static effective barrier ($\alpha = 0.001$), the dynamics resembles that known from fragmented-pair tunneling in a double well [16]. By the Bose-Fermi duality, this may be regarded as two noninteracting fermions Rabi-tunneling independently in the lowest $N_A = 2$ bands $\beta = 0, 1$: $p_R(t) = \sum_\beta \sin^2 \left(J^{(\beta)} t \right) / N_A$. With

increasing α , from our previous discussion we expect an effective attraction between these two fermions. In the intermediate regime ($\alpha = 0.01$), this bears little effect and only leads to a renormalization of the two “Rabi” frequencies $J^{(\beta)}$. However, for mass ratios as large as $\alpha = 0.12$, the picture changes qualitatively: The two atoms tunnel only on a time scale about four times longer than the Rabi oscillations. This pattern closely resembles that of correlated pair tunneling. Indeed, a look into the two-body correlations [Fig. 3(c), inset] reveals that the pair probability p_2 stays remarkably close to unity, in marked contrast with the conventional fragmented-atom pair. In this light, it is enticing to think of this as the tunneling of an attractively *bound pair of identical fermions*.

Let us account for this in a simplified model. Modifying the derivation of (2) for the case of noninteracting *fermions*, we find an effective multi-band Hubbard model [14]. For the special case of $N_A = 2$ “fermions”, this takes the form

$$\bar{H}_A^{(\text{FH})} = - \sum_{\langle s, s' \rangle, \beta} J^{(\beta)} \hat{f}_s^{(\beta)\dagger} \hat{f}_{s'}^{(\beta)} + \delta u \sum_s \hat{n}_s^{(0)} \hat{n}_s^{(1)},$$

where N_A bands $\beta = 0, 1$ contribute; note that we have discarded the on-site term $\sum_{s, \beta} \epsilon^{(\beta)} \hat{n}_s^{(\beta)}$ constant in our setup. The induced on-site interaction turns out to be negative, $\delta u < 0$. For larger α , the induced interaction $|\delta u| \gg J^{(\beta)}$ shifts single-atom tunneling far off resonance. It is in this limit that we can understand the inter-species tunneling as that of an attractively bound fermion(ized) pair at a fairly large tunnel period $T/2\pi \sim |\delta u|/4J^{(0)}J^{(1)}$. From this viewpoint, the two quasi-bound fermions reside in different bands, i.e, pertain to different pseudo-spins and thus are not constrained by Pauli’s principle. This picture extends to arbitrary numbers of fermions $N_A > 2$, which tend to tunnel in dynamically bound N_A -atom clusters.

In conclusion, we have investigated the tunneling of bosonic atoms through a second, localized, species. Remarkably, this can be well understood as an induced *attraction* between the mobile bosons. For the ground state, it may lead to strong intra-species correlations even in the absence of intra-species interactions. The inter-species dynamics, among other things, features tunneling of essentially an attractively bound fermionized pair. Extending the discussion of tunneling to nonclassical potential barriers opens up intriguing perspectives, such as creating an effective lattice made of localized atoms. This would allow for the study of disorder beyond the “quenched” static limit, to give but one example.

Financial support from Landesstiftung Baden-Württemberg and from Leopoldina (S.Z.) is gratefully acknowledged. The authors thank H.-D. Meyer, S. Jochim, T. Gasenzer, and A. Jackson

for valuable discussions.

- [1] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
- [2] S. Fölling *et al.*, *Nature* **448**, 1029 (2007); K. Winkler *et al.*, *Nature* **441**, 853 (2006).
- [3] M. Albiez *et al.*, *Phys. Rev. Lett.* **95**, 010402 (2005); T. Anker *et al.*, *Phys. Rev. Lett.* **94**, 020403 (2005).
- [4] M. Greiner *et al.*, *Nature* **415**, 39 (2002).
- [5] S. Trotzky *et al.*, *Science* **319**, 295 (2008).
- [6] H.-D. Meyer, U. Manthe, and L. S. Cederbaum, *Chem. Phys. Lett.* **165**, 73 (1990); M. H. Beck *et al.*, *Phys. Rep.* **324**, 1 (2000).
- [7] S. Zöllner, H.-D. Meyer, and P. Schmelcher, *Phys. Rev. A* **78**, 013629 (2008).
- [8] E. Haller *et al.*, *Science* **325**, 1224 (2009).
- [9] T. Gericke *et al.*, *Nature Phys.* **4**, 949 (2008).
- [10] M. Girardeau, *J. Math. Phys.* **1**, 516 (1960).
- [11] S. Zöllner, H.-D. Meyer, and P. Schmelcher, *Phys. Rev. A* **74**, 053612 (2006); *Phys. Rev. A* **74**, 063611 (2006).
- [12] O. Mandel *et al.*, *Phys. Rev. Lett.* **91**, 010407 (2003).
- [13] J. Goold and T. Busch, *Phys. Rev. A* **77**, 063601 (2008); X. Yin *et al.*, *Phys. Rev. A* **78**, 013604 (2008).
- [14] A. C. Pflanzner, S. Zöllner, and P. Schmelcher, to be published.
- [15] D. Jaksch *et al.*, *Phys. Rev. Lett.* **81**, 3108 (1998).
- [16] S. Zöllner, H.-D. Meyer, and P. Schmelcher, *Phys. Rev. Lett.* **100**, 040401 (2008).
- [17] R. Piil and K. Mølmer, *Phys. Rev. A* **76**, 023607 (2007); M. Valiente and D. Petrosyan, *J. Phys. B* **41**, 161002 (2008).